

AD-A127 996 FINAL REPORT(U) COLUMBIA UNIV NEW YORK  
D V CHUDNOVSKY ET AL. 1983 N00014-78-C-0318


1/1

UNCLASSIFIED

F/G 20/10

NL

							END DATE FILMED 6 83 DTIC
--	--	--	--	--	--	--	---------------------------------------



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

ONR GRANT N00014-78C-0318

Principal Investigator: Lipman Bers

Grant Period: 3/1/78 - 6/30/82

(12)

Bers

DTIC  
ELECTE  
MAY 10 1983

B

Final Report

DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

D.V. Chudnovsky and G.V. Chudnovsky

*COLUMBIA UNIVERSITY, NEW YORK, N.Y. 10027*

D.V. and G.V. Chudnovsky have successfully continued their studies of internal symmetries of important physical continuous and lattice (discrete) models, and the construction of exact lattice equivalents of continuous systems. These studies were based on the investigation of canonical (Bäcklund) transformations of multidimensional dynamical systems and properties of the symplectic structure of completely integrable systems in the classical and quantum cases.

During the years of the grant, the main subject of the investigation was an analysis of the intrinsic properties of two dimensional classical and quantum, and three dimensional classical physically relevant systems with (hidden) symmetries. The methods of Bäcklund transformations (BTs for short) were shown to play a crucial role in the determination of possible symplectic structure, the determination of their conservation laws and their actual solution [1], [2], [3], [4], [5], [6]. The concept of Bäcklund transformations (BTs) in its most general form was described by the authors as the isomonodromy deformation associated with the Riemann boundary value problem. This description of BTs can be traced back to Riemann, Lappo-Dani'evsky, Garnier, et al. (see

DTIC FILE COPY

ADA 127996

83 03 11 001

the introduction to the investigators' book [7]).

*Bäcklund transformations*  
Using the method of ~~BTs~~ the investigators studied in detail classical and quantum two dimensional completely integrable systems, and lattice models equivalent to and approximating these continuous systems were constructed and examined. Among the continuous systems thoroughly examined in this way are the following important models of quantum field theory and statistical mechanics: the matrix (quantum and classical) nonlinear Schrodinger equation, the sin-Gordon equation, the matrix (non-Abelian) Toda lattice, the generalized XXX-and XXZ-quantum models and models of statistical mechanics, and the quantum generalized XYZ-lattice system (whose classical counterpart is the Landau-Lifshitz equation of superconductivity and whose statistical mechanics reductions include the eight-vertex model of Baxter), [1], [4], [5], [6], [8], [9], [10], [11].

Important properties of Bäcklund transformations, that allow the construction of completely integrable systems and the determination on various symplectic structures associated with them, were described using the language of category theory and algebraic topology [12], [13], [10]. In this way classical and quantum systems were studied simultaneously using the algebra of BTs and the language of factorization axioms for the quantum S-matrices [10], [12], [5], [14].

The investigators applied the algebra of BTs to the three dimensional classical systems as well. Various identities between BTs produce the uniform description of all known matrix two dimensional completely integrable systems as subsystems of scalar three-

dimensional equations of Zakhorov-Shabat type (equivalent to plasma physics equations of Kadomtzev-Petviashvili form) [10], [15]. These studies supplemented the earlier studies of the investigators on the new classes of infinite-dimensional Lie algebras [6], [8], [10].

Methods of BTs were used by the investigators to study Padé approximations of functions satisfying linear differential equations. This was achieved using the interpretation of BTs as isomonodromy transformations and using the monodromy representation of Padé approximations. These connections were discovered for the first time by the investigators [3], [16], [17], [18]. Explicit Padé approximations were studied for algebraic functions and generalizations of Gauss hypergeometric functions. This provides us with new applications to the study of the diophantine approximations of algebraic numbers and values of the logarithmic and inverse trigonometric functions, including cubic irrationalities and numbers such as  $\pi/\sqrt{3}$ , cf. [17].

In 1981 the investigators visited and lectured at the Mittag-Leffler Institute in Stockholm (Sweden) during the special year of Mathematical Physics, the Laboratory of Theoretical Physics at Ecole Normale Supérieure, Paris (France) and the Department of Applied Mathematics Weizmann Institute at Rehovoth (Israel). During the year 1981-82 the investigators were on Guggenheim fellowships.

The support of ONR was important for many of the investigators' projects. In 1982 the volume of proceedings of the investigators' seminar of 1979-80 on the Riemann problem was published [7]. The support of ONR is explicitly acknowledged with gratitude in the introduction.

## References

- [1] D.V. Chudnovsky, G.V. Chudnovsky, Bäcklund transformation for two dimensional nonlinear evolution equations and decomposition theorem, Phys. Lett. A, 82A, 1981, 271-275
- [2] D.V. Chudnovsky, G.V. Chudnovsky, Bäcklund transformations connecting different isospectral deformation equations, J. Math. Phys. 22, 1981, 2518-2522
- [3] G.V. Chudnovsky, An explicit solution of classical and quantum field theory, models and parallel arithmetical problems. A unified approach, Lecture Notes of Invited Address, Annual Meeting of the American Mathematical Society, San Francisco, 1981.
- [4] D.V. Chudnovsky, G.V. Chudnovsky, The relationship between non-Abelian matrix Toda lattice and Ising-type models of statistical mechanics, Phys. Lett. A, 84A, 1981, 353-356.
- [5] D.V. Chudnovsky, G.V. Chudnovsky, Quantum Hamiltonians associated with finite-dimensional Lie algebras and factorized S-matrices, Phys. Rev. Lett. 47, 1981, 1093-1096.
- [6] D.V. Chudnovsky, G.V. Chudnovsky, Bäcklund transformation as a method of decomposition and reproduction of two-dimensional nonlinear systems Phys. Lett. A, 87A, 1982, 325-329.
- [7] D.V. Chudnovsky, G.V. Chudnovsky, The Riemann Problem, Complete Integrability and Arithmetic Applications, Lecture Notes in Mathematics, v. 925, Springer-Verlag, NY, 1982, iv + 373 pp.

- [8] D.V. Chudnovsky, G.V. Chudnovsky, Bäcklund transformations and lattice systems with G-gauge symmetries, Phys. Lett. A., 89A, 1982, pp. 117-122.
- [9] D.V. Chudnovsky, G.V. Chudnovsky, A. Neve, Classical and quantum operator nonlinear Schrödinger equation. I, Lecture Notes Math., v. 925, Springer 1982, pp. 157-177.
- [10] D.V. Chudnovsky, Bäcklund transformations and geometric and complex-analytic background for construction of completely integrable lattice systems, in Problems in Mathematics and Physics, Gordon and Breach, NY, 198 (to appear).
- [11] D.V. Chudnovsky, G.V. Chudnovsky, Hamiltonian structure of isospectral deformation equations, Elliptic curve case, Lecture Notes Math., v. 925, Springer, 1982, pp. 134-146.
- [12] D.V. Chudnovsky, G.V. Chudnovsky, The construction of factorized S-matrices - the relationship between the Baxter model and "Zamolodchikov algebras," Phys. Lett. B, 98B, 1981, 83-88.
- [13] D.V. Chudnovsky, G.V. Chudnovsky, Completely X-symmetric S-matrices corresponding to Theta functions and models of statistical mechanics, Lett. Math. Physics, 5, 1981, 43-51.
- [14] D.V. Chudnovsky, Approximation semi-classique des matrices S factorisées et structure hamiltonienne, C.R. Acad. Sci. Paris, Serie A, v. 291 (1980), A-599-A-602.

- [15] D.V. Chudnovsky, Topological and algebraic structure of linear problems associated with completely integrable systems, Colloquium on group theoretical methods in Physics, Istanbul, 1982, pp. 1-33.
- [16] G.V. Chudnovsky, Rational and Padé approximations to solutions of linear differential equations and the monodromy theory, Lecture Notes in Physics, Springer, 1980, v. 126, 136-169.
- [17] G.V. Chudnovsky, Bäcklund transformations and deformations of linear differential equations with applications to Diophantine approximations, in Problems in Mathematics and Physics, Gordon and Breach, NY, 1983, (to appear).
- [18] D.V. Chudnovsky, G.V. Chudnovsky, Bäcklund transformation for linear differential equations and Padé approximation I, Journal de Mathematique Pures et Appliques, Paris, France, 1981, pp. 1-16.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
<b>PER LETTER</b>	
By	
Distribution/	
Availability Codes	
Dist	Special
<b>A</b>	



DATE  
FILMED  
8